

Quantum Hall effect at low magnetic fields due to electron-electron interaction

In a recent Letter [1] Huckestein showed that at real experimental conditions it is not possible to observe the quantum Hall effect (QHE) at $\omega_c\tau < 1$ predicted by Khmelnitskii [2] ($\omega_c = eB/m$ is the cyclotron frequency and τ is the scattering time). The observation of such a QHE with reentrant Hall plateaus at low magnetic fields would give support for the global phase diagram [3] of the integer QHE as obtained from scaling theory [2]. We would like to point out here that in fact the situation is not so hopeless due to the influence of the electron-electron interaction.

For a bare high-temperature conductivity $\sigma_{xy}^0 = \sigma_0\omega_c\tau/[1 + (\omega_c\tau)^2] \leq \sigma_0/2 < 1.5e^2/h$ with zero-field conductivity $\sigma_0 = ne^2\tau/m < 3e^2/h$, only one QHE plateau with $\sigma_{xy} = e^2/h$ is possible. For $\sigma_0 > 3e^2/h$ additional QHE plateaus appear and, therefore, a reentrant QHE structure can occur for $\omega_c\tau < 1$ [1]. For the quantum values $\sigma_{xy}^0 = ie^2/h$ the localization length

$$\xi \sim l \exp(\pi^2 \sigma_{xx}^0 / 2h^2 / e^4), \quad (1)$$

is huge (l is the elastic mean free path). For $\sigma_{xx}^0 = \sigma_0/[1 + (\omega_c\tau)^2] > \sigma_0/2 > 1.5e^2/h$ with $\omega_c\tau < 1$, $\xi \gtrsim 10^9 l$ exceeds all rational sizes of samples [1]. Therefore, a sample for which the low-field QHE could be observed due to the localization of electrons in a 2D system [2], in fact will be metallic at $\omega_c\tau < 1$ down to a small field B_{IM} . The value of B_{IM} can be estimated from $\pi\xi_0^2 B_{IM} \sim \Phi_0$, where $\Phi_0 = h/e$ is the flux quantum and $\xi_0 \sim l \exp(\pi\sigma_0 h/2e^2)$ is the zero-field localization length which is much smaller than the localization length at a magnetic field larger than B_{IM} . For $\sigma_0 = 3e^2/h$ and $l = 3 \times 10^{-8}$ m the value $B_{IM} \sim 10^{-4}$ T is small.

However, in addition to the localization effects for non-interacting electrons, the electron-electron interaction results in a logarithmic decrease of the density of states at the Fermi level and of the dissipative conductivity [5]

$$\sigma_{xx} = \sigma_{xx,0} - \lambda e^2 / \pi h \ln(T_0/T) \quad (2)$$

where $\lambda < 1$ is the electron-electron interaction constant of order unity. At low enough temperatures a Coulomb gap should occur with localized states near the Fermi level. The localization length should be minimal at the Fermi energy E_F , increases for $|E - E_F|$ increasing [6], and becomes much larger than the sample size outside the gap in accordance with Eq.(1). Since the dissipative conductance vanishes at zero temperature, then according to Laughlin's gauge argument the Hall conductance should be quantized [4]. Therefore, σ_{xy} should tend to a quantum value with temperature decreasing. Since σ_{xy} attains different quantum values depending on the σ_{xy}^0 value, transition regions should exist for which σ_{xx} has a finite value and σ_{xy} is not quantized at zero temperature.

In this scenario there are no levitating extended states between localized states below the Fermi level, which are necessary for the QHE in the case of noninteracting electrons [2,7]. All states below the Coulomb gap down to the mobility edge near the bottom of the band are delocalized. However, the flow diagram [2] and the global phase diagram [3] of the QHE should be qualitatively valid in this case.

From Eq.(2) follows that the temperature at which one could expect the QHE should be of the order of

$$T_1 \sim T_0 \exp(-\pi\sigma_{xx}^0 h / e^2 \lambda) \quad (3)$$

For $\sigma_{xx}^0 = 1.5e^2/h$ and $\lambda = 1, 0.7$ and 0.5 , the temperature $T_1 \approx 1, 0.1$ and 0.01 K, correspondingly, for a reasonable value $T_0 \approx \hbar/k_B\tau \approx 100$ K.

A support of the above presented scenario is given by the recent observation of the QHE in heavily doped n-GaAs layers at $\omega_c\tau > 1$ but with thicknesses of the samples $d > l$, when the bare electron spectrum is continuous like for bulk material [8]. The QHE is observed for σ_{xx}^0 up to $2.6e^2/h$ with $\xi \sim 10^{29}l$ as calculated from Eq.(1). For the explanation of the QHE in this case, the electron-electron interaction has to be taken into account.

Along these lines one could expect a QHE at low magnetic fields in a two-dimensional disordered system with a sufficient large constant of interaction λ and scattering rate $1/\tau$, at experimentally obtainable temperatures. An enhanced spin-splitting will be in favor for such a situation [9].

S. S. Murzin^{1,2} and A. G. M. Jansen²

¹Institute of Solid State Physics RAS, 142432, Chernogolovka, Moscow distr., Russia

²Grenoble High Magnetic Field Laboratory, MPI-FKF and CNRS, B.P.166, F-38042, Grenoble Cedex 9, France

PACS nubers: 73.40.Hm, 71.30.+h, 72.15.Rn, 71.55.Jv

-
- [1] Bodo Huckestein, Phys. Rev. Lett. **84**, 3141 (2000).
 - [2] D.E. Khmel'nitskii, Pis'ma Zh. Eksp. Teor. Fiz. **38**, 454 (1983) [JETP Lett. **38**, 552 (1983)]; Phys. Lett. **106A**, 182 (1984).
 - [3] S. Kivelson, D.Lee, and S. Zang, Phys. Rev. B **46**, 2223 (1992).
 - [4] R. B. Laughlin, Phys. Rev. B **23**, 5632 (1981).
 - [5] P. A. Lee and T. V. Ramakrishnan, Rev. Mod. Phys. **57**, 287 (1985).
 - [6] Gun Sang Jeon, et.al., Phys. Rev. B **59**, 3033 (1999).
 - [7] R. B. Laughlin, Phys. Rev. Lett. **52**, 2304 (1984).
 - [8] S. S. Murzin, A. G. M. Jansen, and P. v. d. Linden, Phys. Rev. Lett. **80**, 2681 (1998); S. S. Murzin, I. Claus, A. G. M. Jansen et. al., Phys. Rev. B **59**, 7330 (1999).
 - [9] S. W. Hwang, H. P. Wei, L. W. Engel et.al., Phys. Rev. B **48**, 11416 (1993).